

1. PERFORMANCE


- 1) Measuring range : 1-20 ppm
- Number of pump strokes 1 (100mℓ)
- 2) Sampling time : 2 minutes/1 pump stroke
- 3) Detectable limit : 0.1 ppm
- 4) Shelf life : 2 years
- 5) Operating temperature : 0 ~ 40 °C
- 6) Reading : Concentration chart method
- 7) Colour change : White → Orange

2. RELATIVE STANDARD DEVIATION

RSD-low : 10% RSD-mid. : 5% RSD-high : 5%

3. CHEMICAL REACTION

o-Toluidine is oxidized and Orthoquinone is produced.






4. CALIBRATION OF THE TUBE

COLOURIMETRY METHOD

5. INTERFERENCE AND CROSS SENSITIVITY

Substance	Interference	ppm	Coexistence
Nitrogen dioxide FIG.1	Similar stain is produced.		Higher readings are given.
Chlorine FIG.2	∕	1	∕
Chlorine dioxide FIG.3	∕		∕



TEMPERATURE CORRECTION TABLE

Chart Readings (ppm)	Correct Concentration (ppm)					
	0 °C (32° F)	10 °C (50° F)	20 °C (68° F)	30 °C (86° F)	40 °C (104° F)	
20	32	24	20	17	14	
15	22	18	15	12	10	
10	15	13	10	8	6	
5	9	7	5	4	3	
3	6	4	3	2	1.5	
1	3	1.5	1	0.8	0.5	

